

NEW

HALEON

THERAFLU

SEVERE COLD RELIEF

NIGHTTIME + + +

SoftCHEWS

Acetaminophen  
Pain Reliever/Fever Reducer  
Diphenhydramine HCl  
Antihistamine/Cough Suppressant

FAST relief ANYWHERE  
fever / cough / headache  
/ body aches / sore throat  
/ runny nose / sneezing  
/ watery eyes

20 CHEWABLE TABLETS  
Chew tablets completely before swallowing



Apple  
Cinnamon  
Flavor

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Distributed by: Haleon, Warren, NJ 07059  
Trademarks are owned by or licensed to the Haleon group of companies.  
Pat. Info [www.productpats.com](http://www.productpats.com)  
©2023 Haleon group of companies or its licensor.

208805

Non Varnish  
Area

57155563



Peel  
Here

### Drug Facts

#### Active ingredients (in each chewable tablet)

Acetaminophen 325 mg..... Pain reliever/Fever reducer  
Diphenhydramine HCl 12.5 mg..... Antihistamine/  
Cough suppressant

#### Purposes

#### Uses

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - sore throat pain
  - headache
  - runny nose
  - sneezing
  - itchy nose or throat
  - itchy, watery eyes due to hay fever
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever

### Drug Facts (continued)

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: 

- skin reddening
- blisters
- rash

 If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

#### Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen

Bleed Limit

Printable Text Area

### Drug Facts (continued)

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin

#### Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

#### Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

#### When using this product

- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

#### Stop use and ask a doctor if

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain or cough gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not use more than directed
- do not take more than 10 chewable tablets in 24 hours unless directed by a doctor. Take every 4 hours, while symptoms persist.

Age	Dose
adults and children 12 years of age and over	take 2 chewable tablets every 4 hours
children under 12 years of age	do not use

#### Other information

- each chewable tablet contains: calcium 10 mg, potassium 10 mg, sodium 7 mg
- store at a controlled room temperature 20-25°C (68-77°F)

**Inactive ingredients** brown iron oxide, calcium gluconate monohydrate, carboxymethylcellulose calcium, carnauba wax, corn syrup solids, crospovidone, FD&C red no. 40, glycerin, mannitol, natural and artificial flavors, pregelatinized starch, simethicone, sodium gluconate, sorbitol, sorbitol solution, sucralose, sucrose, whole dry milk

**Questions or comments?** call 1-855-328-5259

Hinge Area  
No Print