

USE ONLY AS DIRECTED Theraflu Severe Cold Relief Chest Congestion Daytime (continued)

Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

Stop use and ask a doctor if

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain or cough gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not use more than directed
- measure the dose correctly using the enclosed dosing cup
- take every 4 hours in dosing cup provided, while symptoms persist
- do not take more than 5 doses (150 mL) in 24 hours unless directed by a doctor

Age	Dose
adults and children 12 years of age and over	30 mL
children under 12 years of age	do not use

Other information

- each 30 mL contains: potassium 30 mg, sodium 13 mg
- store at controlled room temperature 20°-25°C (68°-77°F)

Inactive ingredients acesulfame potassium, anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, glycerin, maltitol solution, natural and artificial flavors, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments? 1-855-328-5259

Theraflu Severe Cold Relief Nighttime (continued)

Drug Facts (continued)

Inactive ingredients acesulfame potassium, anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, glycerin, maltitol solution, natural and artificial flavors, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?
1-855-328-5259

DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME. DO NOT TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF THE DAYTIME PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

TAMPER-EVIDENT INNER UNIT.
DO NOT USE IF NECKBAND PRINTED WITH “SEALED FOR SAFETY” IS TORN OR MISSING.



SEVERE
COLD RELIEF
VALUE PACK

MULTI-SYMP TOM COLD RELIEF HALEON



2-8.3 FL OZ (245.5 mL) BOTTLES TOTAL 16.6 FL OZ (491 mL) USE ONLY AS DIRECTED

*Temporarily controls cough to help you rest. This is not a sleep aid.

Theraflu Severe Cold Relief Chest Congestion Daytime

Drug Facts

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/ Fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

- Uses**
 - temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - cough due to minor throat and bronchial irritation
 - temporarily reduces fever
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

- Do not use**
 - in a child under 12 years of age
 - if you are allergic to acetaminophen
 - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease

1-855-328-5259

Distributed by: Haleon, Warren, NJ 07059
Trademarks are owned by or licensed to the Haleon group of companies.
©2024 Haleon group of companies or its licensor.
Pat. Info www.productpats.com
See individual boxes for foreign country of origin.