

TheraFlu

Theraflu Daytime Severe Cold Relief (continued)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• do not use more than directed

• take every 4 hours, while symptoms persist. Do not take more than 6 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

• dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.

Drug Facts (continued)

• If using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

• each packet contains: potassium 10 mg, sodium 20 mg
• phenylephedrine contains phenylephedrine 20 mg per packet
• store at controlled room temperature 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients acesulfame potassium, anhydrous citric acid, aspartame, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, maltodextrin, natural and artificial flavors, silicon dioxide, sodium citrate, soy lecithin, sucrose, tribasic calcium phosphate.

Questions or comments?

1-855-328-5259

Theraflu Nighttime Severe Cold Relief

Drug Facts

Active ingredients (in each packet) Purposes

Acetaminophen 500 mg Pain reliever/fever reducer

Diphenhydramine HCl 25 mg Antihistamine/Cough suppressant

Uses

• temporarily relieves these symptoms due to a cold:

- minor aches and pains
- minor sore throat pain
- headache
- sneezing
- itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation
- temporarily relieves your cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

• more than 4,000 mg of acetaminophen in 24 hours

• with other drugs containing acetaminophen

• 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

• skin redness • hives • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash,

nausea, or vomiting consult a doctor promptly.

Drug Facts (continued)

Do not use

• in a child under 12 years of age

• if you are allergic to acetaminophen

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• with any other product containing diphenhydramine, even one used on the skin

• to sedate a child or to make a child sleep

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

• taking sedatives or tranquilizers

• taking the blood thinning drug warfarin

When using this product

• avoid alcoholic drinks

• marked drowsiness may occur

• alcohol, sedatives and tranquilizers may increase drowsiness

• be careful when driving a motor vehicle or operating machinery

• excitability may occur, especially in children

Theraflu Nighttime Severe Cold Relief (continued)

Drug Facts (continued)

Stop use and ask a doctor if

- fever gets worse or lasts more than 3 days
- new symptoms occur
- pain or cough gets worse or lasts more than 7 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed

• take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

• dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
• If using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information • each packet contains: sodium 19 mg

• phenylephedrine: contains phenylephedrine 12.9 mg per packet

• store at controlled room temperature 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients acesulfame potassium, anhydrous citric acid, aspartame, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, maltodextrin, natural and artificial flavors, silicon dioxide, sodium citrate, soy lecithin, sucrose, tribasic calcium phosphate.

Questions or comments? 1-855-328-5259

DO NOT TAKE THE THERAFLU SEVERE COLD RELIEF DAYTIME AND THERAFLU SEVERE COLD RELIEF NIGHTTIME PRODUCTS AT THE SAME TIME. DO NOT TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

TAMPER EVIDENT INNER UNIT
DO NOT USE IF SEALED THERAFLU PACKET IS TORN OR BROKEN

1-855-328-5259

Distributed by: Haleon, Warren, NJ 07059

Trademarks are owned by or licensed to the Haleon group of companies.

©2024 Haleon group of companies or its licensor.

Pat. Info www.productpats.com

214393

SEVERE COLD RELIEF COMBO PACK

MULTI-SYMP TOM COLD RELIEF

HALEON

TheraFlu

SEVERE COLD RELIEF COMBO PACK

6 x DAYTIME

Acetaminophen Pain Reliever / Fever Reducer
Dextromethorphan HBr Cough Suppressant

6 x NIGHTTIME

Acetaminophen Pain Reliever / Fever Reducer
Diphenhydramine HCl Antihistamine / Cough Suppressant

Hot liquid therapy that relieves:

- / Cough / Sore throat pain
- / Head and body ache / Fever
- / Runny nose (Nighttime only)
- / Sneezing (Nighttime only)

6 DAYTIME PACKETS
6 NIGHTTIME PACKETS
12 TOTAL PACKETS

Honey Lemon Flavor

DIETARY CERTIFIED

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE. KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

DO NOT TAKE THE THERAFLU SEVERE COLD RELIEF DAYTIME AND THERAFLU SEVERE COLD RELIEF NIGHTTIME PRODUCTS AT THE SAME TIME. DO NOT TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

DO NOT TAKE A DOSE OF THE SEVERE COLD RELIEF NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF SEVERE COLD RELIEF DAYTIME PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.

Theraflu Daytime Severe Cold Relief

Drug Facts

Active ingredients (in each packet)

Acetaminophen 500 mg Pain reliever/fever reducer

Dextromethorphan HBr 20 mg Cough suppressant

Uses • temporarily relieves these symptoms due to a cold:

- minor aches and pains
- minor sore throat pain
- headache
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

• more than 4,000 mg of acetaminophen in 24 hours

• 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

• skin redness • hives • rash

If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use • in a child under 12 years of age • if you are allergic to acetaminophen

• with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

• liver disease

• cough that occurs with too much phlegm (mucus)

• cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

• taking the blood thinning drug warfarin

Stop use and ask a doctor if

• fever gets worse or lasts more than 3 days

• redness or swelling is present

• new symptoms occur

• pain or cough gets worse or lasts more than 7 days

• cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.