

160 mm
160 mm



**JARABE**  
Acetaminofén 160 mg por 5 mL  
Alivio del dolor y la fiebre

**Lea detenidamente este inserto antes de utilizar Panadol Niños. Consérvelo por si necesita volver a leerlo. Consulte a su profesional de la salud si tiene alguna duda.**

**1. ¿Qué contiene Panadol Niños?** Cada cucharadita (5 mL) contiene: Acetaminofén 160 mg. La solución también contiene sacarina sódica, saborizante, ácido benzoico, colorante rojo #40.

**Dosis / Modo de empleo**

**2. ¿Cómo debe ser usado Panadol Niños?** Para uso oral únicamente. Utilice siempre la copa dosificadora proporcionada en el empaque. La copa dosificadora tiene varias escalas, refiérase al peso del niño en la tabla para suministrar la dosis adecuada. Mida la cantidad exacta de líquido en la copa dosificadora.

Peso	Edad	Dosis
13-15 kg	2-3 años	6 mL
16-19 kg	4-5 años	7.5 mL
20-25 kg	6-7 años	9 mL
26-31 kg	8-9 años	12 mL
32-42 kg	10-12 años	15 mL

Dosis máxima diaria: 60 mg / kg administrados en dosis divididas de 10 a 15 mg / kg durante el periodo de 24 horas. Se puede repetir cada 6 horas si los síntomas persisten, sin exceder 4 tomas en un periodo de 24 horas, ni por más de 3 días consecutivos a niños sin consultar al médico. Este medicamento produce un alivio temporal de los síntomas, si no se obtiene dicha mejoría después de 48 horas consulte con el médico. No tomar junto con otros medicamentos de prescripción o sin prescripción que contengan acetaminofén o indicados para el alivio del dolor, fiebre, resfrios o para conciliar el sueño. No se recomienda en niños menores de 2 años. Los rangos de edad en la tabla de dosificación son solamente orientativos. Utilice siempre la menor dosis efectiva para tratar los síntomas y en el menor periodo de tiempo necesario.

**Contraindicaciones**

**3. ¿Cuándo no debe usarse Panadol Niños?** No utilice el medicamento si es alérgico o hipersensible al acetaminofén o a cualquier componente de la formulación.

**Advertencias y precauciones**

**4. ¿Qué precauciones deben tomarse cuando se usa Panadol Niños?** Contiene acetaminofén. No utilizar con otros productos que contengan acetaminofén porque puede ocurrir una sobredosis. Nunca suministre más medicamento de la que se muestra en la tabla. No sobrelene la copa dosificadora. La dosis excesiva del acetaminofén puede causar daño grave al hígado. **Antes de tomar o cuando suministre este medicamento consulte a su médico si:** El dolor y la fiebre persisten por más de 3 días, empeora u ocurren nuevos síntomas. Si ha sido diagnosticado con mal funcionamiento del hígado o de los riñones. Está bajo de peso o desnutrido. Está tomando warfarina u otros medicamentos anticoagulantes. Si tiene infección severa, está severamente desnutrido o severamente bajo de peso, lo que puede incrementar el riesgo de acidosis metabólica, cuyos síntomas incluyen: respiración agitada y con dificultad, náuseas, vómito, pérdida del apetito. Consulte a su médico si presenta la combinación de esos síntomas. Lea y siga las instrucciones del empaque. Contiene colorante rojo #40 puede provocar reacciones alérgicas.

**Interacciones**

**5. ¿Está tomando otros medicamentos?** Consulte a su médico si usted/su niño toma warfarina o medicamentos similares, antibióticos (cloranfenicol), tratamiento con estrógenos, antiepilépticos (lamotrigina, fenitoína, fenobarbital, metilfenobarbital, primidona, carbamazepina), antituberculosos (isoniazida, rifampicina), barbitúricos (sedantes, anticonvulsivantes), carbón activado, colestiramina (baja el colesterol), diuréticos (furosemida), antigotosos (probenecid, sulfpirazona), anticolinérgicos o antiespasmódicos, metoclopramida, domperidona, propranolol, zidovudina.

**Reacciones adversas**

**6. ¿Qué reacciones secundarias puede causar Panadol Niños?** Deje de administrar este medicamento y consulte a su médico inmediatamente si:

- Experimenta una reacción alérgica como salpullido en la piel, picazón, con o sin dificultad para respirar, inflamación de labios, lengua, garganta y cara.
- Experimenta descamación de la piel y úlceras bucales.
- Si ha experimentado problemas para respirar luego de tomar ácido acetilsalicílico o antiinflamatorios no esteroideos y experimenta esas reacciones con este medicamento.
- Experimenta reacciones en piel como el Síndrome de Steven Johnson, Necrólisis Epidérmica Tóxica y Pustulosis Exantemática Generalizada Aguda.
- Experimenta sangrado o hematomas inexplicables. Estas reacciones son raras.

**Manejo de sobredosis**

Si el niño toma más cantidad de medicamento de lo indicado, consulte inmediatamente al médico, aunque no presente síntomas, debido al riesgo de daño hepático. La sobredosis de acetaminofén puede causar insuficiencia hepática que puede llevar a un trasplante de hígado o la muerte.

**Información adicional**

- Consérvese a menos de 30°C.
- Manténgase fuera del alcance y vista de los niños.
- Uso pediátrico.
- Presentaciones: Frasco con 30 mL, 60 mL, y 90 mL. Algunas presentaciones no están a la venta en todos los países.

Para mayor información, reporte de situaciones desfavorables con el producto o quejas, contacte a GSK al correo electrónico [fv.aecaricam@gsk.com](mailto:fv.aecaricam@gsk.com) o a los teléfonos:

Costa Rica: (506) 4055-4242 El Salvador: (503) 2136-7801  
Guatemala: (502) 2376-1930 Honduras: (504) 2540-1638  
Nicaragua: (505) 27222215 Panamá: (507) 833-6768



49.5 mm  
PHARMA CODE REF. IS: 424

MASS D 0,5 mm  
MASS E 1,5 mm  
MASS G 1,0 mm

Example Technical information control code

V1 18 - DEC - 2020		champs <small>Consumer Healthcare Artwork Management &amp; Performance Solution</small>		Artwork Information Panel				gsk																		
D2A	<input type="checkbox"/>	Non Production	<input type="checkbox"/>	Production	<input checked="" type="checkbox"/>	SGS No: 6257347	CHAMPS No: 000004197																			
Total Number of Colours including Varnish & Foils: 1						Manufacturing Site: Pharmetique	Site Component No: 10002105																			
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 16.6%;"></td> </tr> <tr> <td style="text-align: center;">1. Black</td> <td style="text-align: center;">2.</td> <td style="text-align: center;">3.</td> <td style="text-align: center;">4.</td> <td style="text-align: center;">5.</td> <td style="text-align: center;">6.</td> </tr> <tr> <td style="text-align: center;">7.</td> <td style="text-align: center;">8.</td> <td style="text-align: center;">9.</td> <td style="text-align: center;">10.</td> <td style="text-align: center;">11.</td> <td style="text-align: center;">12.</td> </tr> </table>												1. Black	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	12.	Approving Market: Caricam-CCA	Site Change Control No: M00899	
1. Black	2.	3.	4.	5.	6.																					
7.	8.	9.	10.	11.	12.																					
Total Special Finishes: 0						Product Market Trade Name: PANADOL CHILDREN	Component Type: Leaflet																			
						Technical Drawing No: PLANO 160x160 mm Ins LQ1	Material Spec No: N/A																			
						Print Process: Offset- Lithography	Pharma Code No: 424																			
						Barcode No:	Bar Width Reduction: 0 Micron																			
						Formulation Code: N/A	Body Text Size: 6pt Leading: 6.5pt																			
						Micro Text: N	Horizontal Scale: 100% Smallest Text Size: 6pt																			
Studio Location: Mexico City, MX																										

AIP\_CHAMPS\_2019\_Version 1

Page 1 of 2

160 mm
160 mm



**Panadol Children**

**SYRUP**

Acetaminophen 160 mg per 5 mL  
Relief of pain and fever

**Interactions**

**5. Are you taking any other medications?** Consult your doctor if you or your child takes warfarin or similar medicines, antibiotics (chloramphenicol), treatment with estrogens, antiepileptic drugs (lamotrigine, phenytoin, phenobarbital, methylphenobarbital, primidone, carbamazepine), anti-tuberculosis drugs (isoniazid, rifampin), barbiturates (sedatives, anticonvulsants), activated charcoal, cholestyramine (low cholesterol), diuretics (furosemide), anti-gout (probenecid, sulfapyrazone), anticholinergic or antispasmodic, metoclopramide, domperidone, propranolol, zidovudine.

**Adverse reactions**

**6. What side effects could Panadol Children cause?** Stop giving this medicine and tell your doctor immediately if: • Experiences allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face. • Experiences skin rash or peeling, or mouth ulcers. • Has previously experienced breathing problems with acetylsalicylic acid or non-steroidal anti-inflammatory drugs and experience a similar reaction with this medicine. • You experience skin reactions like Stevens- Johnson Syndrome, Toxic Epidermal Necrolysis or Acute Generalized Exanthematic Pustulosis. • Experiences unexplained bruising or bleeding. These reactions are rare.

**Overdose**

If your child takes more of the medicine than he or she should, seek medical advice from a doctor immediately even if your child doesn't have any symptoms because of the risk of liver failure. Acetaminophen overdose may cause liver failure which may require liver transplant or lead to death.

**Other information**

- Store below 30°C. • Keep out of sight and reach of children. • Pediatric use.
- Presentations: Bottle with 30 mL, 60 mL and 90 mL. Some presentations are not for sale in all countries.

**Fabricado por/ Manufactured by**

Pharmetique S.A., Colombia  
Titular / Holder: GlaxoSmithKline Costa Rica S.A., Costa Rica.

For more information, report adverse events or product complaints contact GSK by email at [fv.aecaricam@gsk.com](mailto:fv.aecaricam@gsk.com) or by phone:  
Jamaica: +1 (876) 630 2074  
Trinidad & Tobago: +1 (868) 224 5759

Fecha de la última revisión: Marzo 2019, basado en el GDS v7.0 de Agosto 2017 / Date of the last revision: March 2019, based on the GDS v7.0 of August 2017

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424


49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm



PHARMA CODE REF. IS: 424

PHARMA CODE REF. IS: 424

49.5 mm