

# Infants' Advil

Concentrated Drops

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Child-Resistant Safety Cap

### Drug Facts (continued)

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

#### Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- stomach bleeding warning applies to the child
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has a history of stomach problems, such as heartburn
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child is taking a diuretic

#### Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs

#### Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.

### Drug Facts (continued)

- repeat dose every **6-8 hours**, if needed
- do not use more than **4 times a day**
- measure with the dosing device provided. Do not use with any other device.

#### Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lb	6-11 mos	1.25 mL
18-23 lb	12-23 mos	1.875 mL

#### Other information

- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

#### Inactive ingredients

acetic acid, artificial flavor, butylated hydroxytoluene, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

#### Questions or comments?

Call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

Rinse the oral syringe with tap water only.



# Infants' Advil

Concentrated Drops

## Ibuprofen Oral Suspension

50 mg per 1.25 mL

### Pain Reliever/Fever Reducer (NSAID)

**Fever**

Aches & Pains

Lasts Up To

# 8

Hours

## Dye-Free

White Grape Flavored Drops  
for ages 6-23 months

1/2 FL OZ (15 mL)

Alcohol-Free

**Important:** Read all product information before using. Keep this box for important information. This product is intended for use in children ages 6 months - 23 months.

### Drug Facts

#### Active ingredient Purpose (in each 1.25 mL)

Ibuprofen 50 mg Fever reducer/  
(NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

#### Uses temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, headaches and toothaches

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Dist. by: GSK CH, Warren, NJ 07059

For most recent product information, visit [www.Advil.com](http://www.Advil.com).

©2022 GSK or licensor

000079546