

TOP PANEL 1

NO COPY 1/16" (0.0625") FROM DIE. DOTTED LINE FOR HOLDING ONLY AND DOES NOT PRINT.

KISS CUT

ART BLEEDS OFF 1/16" (0.0625") OUT FROM DIE. DOTTED LINE FOR HOLDING ONLY AND DOES NOT PRINT.

THIS LINE INDICATES DIE AND DOES NOT PRINT.

3/16" X 3/16" DATA MATRIX CODE (ECC 200), PRINTED IN BLACK ON UNPRINTED WHITE BACKGROUND.

1" FOR LOT & EXPIRATION AREA. UNVARNISHED AND PRINTED IN BLACK.

Easy Open ARTHRITIS CAP

HALEON

EASE OF USE CERTIFIED

Arthritis Foundation

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Drug Facts

Active ingredient (in each capsule) Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*Pain reliever/Fever reducer (present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

213112

FPO

160 Liquid Filled Capsules

Solubilized Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)

Actual Size

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

1/8" (0.125") CORNER RADIUS TYP

6.75"

7.25"

0.5"

2.625"

Process CMYK

Pantone 302 C

High Gloss Varnish

BASE PANEL 3

3/8" HINGE AREA UNVARNISHED (LINES DO NOT PRINT)

ART BLEEDS OFF 1/16" (0.0625") OUT FROM DIE. DOTTED LINE FOR HOLDING ONLY AND DOES NOT PRINT.

THIS LINE INDICATES DIE AND DOES NOT PRINT.

NO COPY 1/16" (0.0625") FROM DIE. DOTTED LINE FOR HOLDING ONLY AND DOES NOT PRINT.

NO RELEASE COATING AREA

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients
FD&C green no. 3, gelatin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

Questions or comments? call toll free 1-800-88-ADVIL

For most recent product information visit www.Advil.com

Distributed by: Haleon, Warren, NJ 07059

Trademarks are owned by or licensed to the Haleon group of companies.

©2024 Haleon group of companies or its licensor.

LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.

1/8" (0.125") CORNER RADIUS TYP

6.75"

7.25"

0.5"

2.625"

Base Pantone 302 C

RELEASED BACK PANEL 2

3/8" HINGE AREA UNVARNISHED (LINES DO NOT PRINT)

ART BLEEDS OFF 1/16" (0.0625") OUT FROM DIE. DOTTED LINE FOR HOLDING ONLY AND DOES NOT PRINT.

THIS LINE INDICATES DIE AND DOES NOT PRINT.

NO COPY 1/16" (0.0625") FROM DIE. DOTTED LINE FOR HOLDING ONLY AND DOES NOT PRINT.

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

1/8" (0.125") CORNER RADIUS TYP

6.75"

2.625"

Back Pantone 302 C

V2
17 - JUN - 2024

Artwork Information Panel

HALEON

Non Production Artwork <input type="checkbox"/>	Production Artwork <input checked="" type="checkbox"/>	Component No: 213112	AWP No: LKH
Total Number of Colours including Varnish & Foils: 8		Manufacturing Site: N/A <small>(Note: North American Artwork will have N/A in this field)</small>	
		Approving Market: USA	
1. Process Cyan	2. Process Magenta	3. Process Yellow	4. Process Black
		5. Front Top Ply Pantone 302 C	6. Back Top Ply Pantone 302 C
7. Base Pantone 302 C	8. High Gloss Varnish		
Total Special Finishes: 0		Technical Drawing No: DRW-2696	
		Material Spec No: N/A <small>(Note: North American Artwork will have N/A in this field)</small>	
		Body Text Size: Smallest Text Size: <small>(Note: For North American Artwork, see Regulatory Spec box for this information)</small>	
		Electronic Verification Code: 213112, Color = Process Black	
		Printer and User notes: PRINTER TO REPLACE ALL FPO (For Position Only) CODES WITH SCANNABLE ART. PRINTER RESPONSIBLE FOR ALL TRAPPING, BLEEDS, COATINGS AND RESOLUTION. PRINTER AND USERS RESPONSIBLE FOR PROPER LICENSING OF ALL FONTS USED IN THIS ARTWORK.	

REGULATORY LEGEND

	Actual Size
DRUG FACTS TITLE	9 pt
DRUG FACTS CONTINUED	8 pt
HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6 pt
LEADING	6.5 pt
# OF CHARACTERS PER INCH	<39
BULLETS	5 pt
SPACE BEFORE BULLET	2 ems
BARLINES, HAIRLINES	1.5 pt, .5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 ems
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	13.23 pt
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A