





COLD+FLU
COLD & FLU NEW FORMULATION

**FOR EFFECTIVE RELIEF OF COLD & FLU SYMPTOMS, ACHEs AND FEVER.
INCLUDES EXTRA VITAMIN C. NON-DROWSY FORMULA.**

Description
Each PANADOL Cold & Flu New Formulation tablet contains Acetaminophen (Paracetamol) 500mg, Caffeine 25mg, Phenylephrine Hydrochloride 5mg, Vitamin C 30mg and Terpin Hydrate 20mg.

Excipients
Povidone, purified talc, starch maize, potassium sorbate, sodium lauryl sulphate, stearic acid, eurocod sunset yellow, starch pregelatinized

Actions
PANADOL Cold & Flu tablets contain multi-ingredient formula that could relieve symptoms of cold and flu fast. Paracetamol produces analgesia by elevation of the pain threshold and antipyresis through action on the hypothalamic heat-regulating center, which does not irritate stomach. Phenylephrine Hydrochloride provides fast relief of nasal congestion with long lasting efficacy. Caffeine is a stimulant to relieve the tiredness caused by cold without drowsiness. Wet patient feel comfortable. Vitamin C could help patient recover fast during cold period. Terpin hydrate is an effective expectorant by increasing bronchial secretion.

Indications
PANADOL Cold & Flu tablets provide effective temporary relief of the symptoms associated with common cold and other upper respiratory infections including nasal congestion, sore throat, fever, chesty cough, headache, body ache.

Contraindications
Do not use if:
• you are allergic to paracetamol, ascorbic acid, caffeine, terpin hydrate, phenylephrine or any of the other ingredients in the products.
• you are taking, or have taken in the past two weeks, drugs called monoamine oxidase inhibitors (MAOIs), usually used to treat depression.
• you are pregnant.

Dosage / Usage
Adults and children (12 years and over): 2 tablets every 4 hours or above as required. Do not take more than 8 tablets in 24 hours. Do not exceed the stated dose and do not take more frequently than every 4 hours. Do not use for longer than 7 days without asking your doctor. Always use the lowest effective dose to relieve your symptoms.
Children under 12 years: Do not use in children under 12 years of age.

Warnings and Precautions
• Do not take with any other medicines for the relief of colds and flu, congestion or blocked nose.
• This product contains paracetamol. Taking too much paracetamol can cause serious harm to your liver.
• Do not use this medicine if you are taking any other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu, or to aid sleep.
• Always read and follow the label.
Check with your doctor before use if you:
• have liver or kidney problems.
• are underweight or malnourished.
• regularly drink alcohol.
You may need to avoid using this product altogether or limit the amount of paracetamol that you take.
Check with your doctor before use if you:
• have a severe infection as this may increase the risk of metabolic acidosis.
Signs of metabolic acidosis include:
• deep, rapid, difficult breathing
• feeling sick (nausea), being sick (vomiting)
• loss of appetite
Contact a doctor immediately if you get a combination of these symptoms.
• have high blood pressure, heart disease, blood vessel disease such as Raynaud's phenomenon (which may appear as pain in the fingers or toes in response to cold or stress), diabetes, an overactive thyroid,

glaucoma (excessive pressure inside your eyes), rheochromocytoma (a tumour near the kidney), or prostate problems.
• are taking medicines to control your blood pressure, such as beta-blockers.
• are taking appetite suppressants or stimulants
• are taking medicines to treat depression, such as tricyclic antidepressants (e.g. amitriptyline).
• are taking medicines for heart disease (e.g. digoxin).
• are taking warfarin or similar medicines used to thin the blood.
• are breast-feeding. Use during breast-feeding should be avoided. This medicine contains caffeine. Avoid drinking too many caffeine containing drinks (e.g. tea, coffee and caffeine-containing canned drinks) when taking this medicine.
Please see your doctor if your symptoms persist.
Keep out of sight and reach of children.
This medicine may cause dizziness. If affected do not drive or operate machinery.

Adverse reactions
When using this medicine you may experience: headache, dizziness, nervousness, difficulty sleeping, increased blood pressure, nausea and vomiting.
Stop taking this medicine and tell your doctor immediately if:
• you experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
• you experience a skin rash or peeling, or mouth ulcers.
• you have previously experienced breathing problems with aspirin or non-steroidal anti-inflammatory, and experience a similar reaction with this product.
• you experience unexplained bruising or bleeding.
• you experience loss of vision, which may be due to abnormally high blood pressure in the eye. This is very rare but is more likely to occur in those with glaucoma.
• you experience an unusually fast pulse rate, or a sensation of an unusually fast or irregular heartbeat
• you experience difficulty in passing water. This is more likely to occur in men with an enlarged prostate gland.
These reactions are rare.

Overdose
If you take more of the medicine than you should:
seek medical advice immediately even if you do not have any symptoms because of the risk of liver failure.

Storage Condition
Store below 30°C. Store in a dry place.
Last revision: Dec, 2017
Hong Kong
© 2017 GSK group of companies or its licensor.
Trade marks are owned by or licensed to the GSK group of companies.

CCX-907(HK)
919214



PHARMA CODE REF. IS: 1349





傷風+感冒
傷風感冒丸新配方

**主治：傷風感冒引起的各種症狀、發燒及身體痛楚。
不會使人昏昏欲睡。含更多維他命C。**

成份
「必理痛」傷風感冒丸新配方每粒含有 Acetaminophen (Paracetamol) 500mg, Phenylephrine Hydrochloride 5mg, Caffeine 25mg, Vitamin C 30mg, Terpin Hydrate 20mg

賦形劑
Povidone, purified talc, starch maize, potassium sorbate, sodium lauryl sulphate, stearic acid, eurocod sunset yellow, starch pregelatinised.

作用
「必理痛」傷風感冒丸由於所含的各成份是以適當的比例，精心調配而成，故能迅速舒緩感冒、流行性感冒和其他呼吸系統感染所引起的各種症狀。Paracetamol是一種劑效性極佳的止痛劑，不會刺激胃部，對消除中度疼痛特別有效；亦可消除發熱和由感冒引起的其他痛楚。它的解熱性質亦可降低由呼吸系統感染所引起的發燒。Phenylephrine Hydrochloride是一種耐受性極佳，迅速而長效的鼻充血消除劑，能迅速治療感冒所引起的鼻塞，收縮充血之鼻黏膜，促進鼻道的暢通，恢復舒適的呼吸。Caffeine是一種興奮劑，協助消除感冒所引起的疲勞，促進清醒，使您倍感舒適。維他命C能促進感冒迅速痊癒。Terpin Hydrate能增加支氣管分泌，是一種有效祛痰劑。

療效
舒緩傷風感冒的各種症狀：鼻塞、咽喉痛、發燒、痰多咳嗽、頭痛、身體痛楚。

禁忌症
如有以下情況，切勿服用此藥：
• 對此藥物內任何成份敏感人士。
• 如正服用或在過去兩週期間曾服用單胺氧化酶抑制劑的人士。
• 如您正在懷孕。

劑量和使用法
成人及兒童（十二歲或以上）：每次服兩粒，需要時每隔四小時或以上服一次。二十四小時內服用不超過八粒。不可服用多於指定份量。
請勿服用少於四小時服用此藥物。
如沒有醫生指示，切勿連續服用此藥超過7日。
請使用最低有效劑量，以減輕症狀。
不建議十二歲以下兒童服用此藥物。

注意事項
• 服用此藥時請避免使用其他過熱藥或舒緩傷風感冒的藥物。
• 此產品含有撲熱息痛。服用過量撲熱息痛可導致肝臟受損。
• 如正服用其他處方或非處方含撲熱息痛藥物以治療痛楚、發燒、傷風感冒症狀或幫助睡眠，請勿同時服用此藥。
• 請仔細閱讀並從指示服用「必理痛」傷風感冒丸。
請以下人士在服用此藥前向醫生查詢：
• 患有肝臟或腎臟疾病。
• 您的體重偏輕或營養不良。
• 您有飲酒的习惯。
您可能需要避免服用此產品或減低撲熱息痛的劑量。
如有以下狀況，在服用此藥前請先諮詢醫生：
• 您有嚴重感染，嚴重感染會增加代謝性中毒的風險。代謝性中毒的症狀包括：
• 呼吸深、急促及困難
• 感到不適（噁心）；身體不適（嘔吐）
• 食慾不振
如出現多於一個以上的症狀，請立即求醫。
• 患有高血壓、心臟病、心血管病(如 Raynaud's Phenomenon) —當遇到寒冷或壓力的時候，手指或腳尖可能會感到疼痛)、糖尿病、過度活躍甲狀腺、青光眼(眼睛內部壓力過大)、嗜睡(眼睛腫(腎上腺髓質腫)或前列腺問題)
• 正服用降血壓藥如 beta-blocker。
• 正服用增加或阻塞胃口的藥物。
• 正服用抗抑鬱的藥物如 amitriptyline(其中一種三環抗抑鬱藥)。
• 正服用心臟病藥如 digoxin。
• 正服用 Warfarin或其他抗凝血藥。
• 正在授乳。在授乳期間應避免服用此藥。
此藥物含有咖啡因，在服用本藥物期間應避免飲用過多含咖啡因的飲品(如茶、咖啡、含咖啡因的罐裝飲品)。服用後若症狀持續，請向醫生查詢。
請把藥物貯放在兒童視線以外及不易觸及的地方。服用後如有量感症狀，請勿駕駛或操作機器。

不良反應
服用此藥後，您可能會出現以下情況：頭痛、暈眩、焦慮、難以入睡、血壓上升、噁心及嘔吐。
如服此藥後出現以下不良反應，請立即停藥並延醫診治：
• 出現敏感症狀，如皮膚出疹或瘙癢，間中還出現呼吸困難、喉嚨、舌頭、咽喉或面部腫脹。
• 皮膚出疹、脫皮或口腔出現潰瘍。
• 曾因服用阿士匹靈或其他非類固醇類消炎藥出現呼吸困難的症狀，及在服用本藥物期間出現相似症狀。
• 出現不明原因或流血。
• 失明(可能因眼內出現異常高血壓所致)，這是十分罕見的，但青光眼患者較易有此症狀。
• 不尋常的快速脈搏率又或者感到不尋常快或無規律的心跳。
• 排尿困難。患有前列腺肥大的男性服用此藥較易有此症狀。
這些不良反應是罕見的。

服藥過量
若服用過量撲熱息痛，可導致肝衰竭。因此即使用過量藥物後並無任何症狀，亦應儘速治療。

貯存方法
貯存於30°C以下及乾爽處。
Last Revised: Dec, 2017

CCX-907(HK)
919214



PHARMA CODE REF. IS: 1349

SGS Number: 733402		Item Number: 919214		Manufacturing Site: GSK-MYS-Kuala Lumpur-MYKLU	
GlaxoSmithKline Artwork Information Panel		Market Trade Name: Panadol Cold & Flu		Colour Standard Reference Number: N/A	
Market or Pack Owner: CH-Hong Kong-HKG		Printing Process: Offset		Substrate: White paper	
Technical Reference No(s).: IMA_C80_INS_131X208_DRW (do NOT include the technical reference doc(s) version no(s).)					
Colours	348	Varnishes		Special Finishes	
Total: 1		Total: N/A		Total: N/A	
				RSC A/W Version: 6	

Artwork copyright is the property of the GlaxoSmithKline Group of Companies

All suppliers providing a service to GSK for printed components of any description must ensure that they have a licence for all fonts / software used in conjunction with GSK artwork. The distribution and use of fonts / software without a licence constitutes an intellectual property infringement. GSK will not accept any liability for the breach of third party intellectual property rights by printed component suppliers. The GSK certification / audit process requires suppliers to declare that they do not use unlicensed fonts / software and may require the supplier to produce evidence of such licence to GSK.

ATTENTION • ATTENTION • ATTENTION • ATTENTION • ATTENTION • ATTENTION • ATTENTION • ATTENTION

To Ensure Accurate PDF Viewing and Printing:
FOR SCREEN VIEWING: Use Adobe Acrobat 7 Professional or Adobe Acrobat Reader, Standard or Professional (higher than 7).
Overprint Preview must be activated for accurate on screen viewing.
FOR PRINTING: Use only Acrobat Professional version 7 or higher. "Apply Overprint Preview" or "Simulate Overprinting" must be activated in the print settings for printing accurate hard copies.

180 mm Measuring Bar

If a status identification banner DOES NOT appear on this document, THEN this document has NOT been printed from the Global Pack Management system.

AP_Disclaimers_A1_SEP_2013 Version 1

IMA_C80_INS_131X208_DRW

2

IMPORTANT

GSK Market is responsible for this product, its design and content.
Ensure the artwork is thoroughly checked, all the text proof-read and approved.
RSC GSK is responsible for site technical requirements and pre-press suitability.

GSK Market is responsible to advise RSC in case changes required impact the followings:

Formulation
Tablet embossing
Storage conditions
Shelf Life

FONT REPLACED

Market to check all text 100%

TEXT SIZE CONTAINED IN THIS ARTWORK

Body text size: 5.0pt

Leading: 7.3pt

Horizontal Scale: 100%

Smallest text size: 4.5pt

Microtext: N