 e-Banner	Project: CO-0002771	Document: PPC-1500163	Version: 6
	Site Code: 910554	Operator: ED769180	Date/Time Created: 14.Dec.2018 10:54 GMT

FRONT





COLD+FLU
COLD & FLU EXTRA

FOR EFFECTIVE EXTRA RELIEF OF COLD & FLU SYMPTOMS: ACHES AND FEVER. INCLUDES EXTRA VITAMIN C. NON-DROWSY FORMULA.

Description:
Each Panadol Cold & Flu Extra Caplet contains:
Acetaminophen (Paracetamol) 500mg, Phenylephrine Hydrochloride 10mg, Noscapine 15mg, Caffeine 25mg, Ascorbic Acid 63.33mg, Terpin Hydrate 33.33mg.

Dosage:
For Oral Administration only.
Adults and children (over 12 years): Take 2 caplets every 6 hours, do not take more than 8 caplets in 24 hours. Do not take more frequent than every 6 hours. Do not use for longer than 7 days without asking your doctor.
Should not be used with other Acetaminophen containing products, decongestants, or cold and flu medicines. Do not exceed the stated dose.
Children 12 years and under: Not recommended for children of 12 years and under.

Actions:
Panadol Cold & Flu Extra caplet contains multi-ingredient formula that could relieve cold and flu symptoms fast. Acetaminophen produces analgesia by elevation of the pain threshold and antipyresis through action on the hypothalamic heat-regulating center, which does not irritate stomach. Phenylephrine Hydrochloride has fast relief of nasal congestion with long lasting efficacy. Caffeine is a stimulant to relieve the tiredness caused by cold without drowsiness, let patient feel comfortable. Vitamin C could help patient to recover fast during cold period. Terpin Hydrate is an effective expectorant by increasing bronchial secretion. Noscapine is a centrally acting cough suppressant which is used to relieve non-productive cough.

Indications:
Panadol Cold & Flu Extra Caplets provide effective temporary relief of the symptoms associated with common cold including fever, aches, nasal and sinus congestion, cough and minor throat pain, headache.

Contraindications:
Panadol Cold & Flu Extra Caplets is contraindicated in patients with a previous history of hypersensitivity to any of the ingredients. This product is also contraindicated in patients who are taking or have taken within the last two weeks, monoamine oxidase inhibitors. Do not use if you are pregnant.

Precautions:
1. If you are taking other analgesics, antipyretics or cold relief medicine, don't take this medicine.
2. Please see your doctor if symptoms persist.
3. Keep this and all medication out of reach of children.
4. **Please check with your doctor before use if:**
a. You are taking Warfarin or similar medicines used to thin the blood (The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of Acetaminophen with increased risk of bleeding; occasional doses have no significant effect).
b. You have high blood pressure, heart disease, blood vessel disease such as Raynaud's phenomenon (which may appear as pain in the fingers or toes in response to cold or stress), diabetes, an overactive thyroid, angle closure glaucoma, pheochromocytoma (a tumour near the kidney), or prostate problems.
c. You are taking medicines to control your blood pressure such as beta-blockers.
d. You are taking appetite suppressants or stimulants.
e. You are taking medicines to treat depression, such as tricyclic antidepressants (e.g. amitriptyline).
f. You are taking medicines for heart disease (e.g. digoxin).
g. You have been diagnosed with liver or kidney impairment.
h. You are breast-feeding. Use during breast-feeding should be avoided.

Adverse reactions:
Toxicity is extremely rare following administration of recommended dose. Nevertheless, massive overdose of acetaminophen may adversely affect the liver of some people and signs or symptoms of overdose may be delayed for up to one week following ingestion. Nevertheless, liver damage is largely reversible if the patient is treated early, thus **immediately medical management is required in the event of overdose, even if symptoms of overdose are not present.** Acetaminophen overdose may cause liver failure.

Adverse reactions:
Undesirable effects are rare. When using this medicine you may experience: headache, dizziness, nervousness, difficulty sleeping, increased blood pressure, nausea and vomiting.

Stop taking this medicine and tell your doctor immediately if you have:
a. Allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
b. A skin rash or peeling, or mouth ulcers.
c. Previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction with this product.
d. Unexplained bruising or bleeding.
e. Loss of vision, which may be due to abnormally high blood pressure in the eye. This is very rare but is more likely to occur in those with glaucoma.
f. An unusually fast pulse rate, or a sensation of an unusually fast or irregular heartbeat.
g. Difficulty in passing water. This is more likely to occur in men with an enlarged prostate gland.

Last revision: Oct 2018
Hong Kong

© 2018 GSK group of companies. All rights reserved.
Trade marks are owned by or licensed to the GSK group of companies.

CDP-904(HK)
910554

PHARMA CODE REF: IS-1358

BACK





傷風+感冒
傷風感冒特強

主治：更有效舒緩傷風感冒引起的症狀：發燒及身體痛楚。不會使人昏昏欲睡。含更多維他命C。

成份
「必理痛」傷風感冒特強每粒含有Acetaminophen (Paracetamol) 500mg, Phenylephrine Hydrochloride 10mg, Noscapine 15mg, Caffeine 25mg, Ascorbic Acid 63.33mg, Terpin Hydrate 33.33mg。

服法
只供口服
成人及兒童 (十二歲以上)：每次服兩粒，每隔六小時服用一次，二十四小時內服用不超過八粒。請勿於6小時內重複服用。如沒有醫生指示，切勿連續服用此藥超過7日。
服用時請勿服用其他含Acetaminophen、通鼻塞藥或其他傷風感冒的藥物。不可服用多於指定份量。
十二歲或以下兒童：不建議十二歲或以下兒童服用此藥物。

作用
「必理痛」傷風感冒特強由於所含的各成份是以適當的比例，精心調配而成，故能迅速舒緩感冒引起的各種症狀；Acetaminophen能提升對痛楚的忍耐力，從而舒緩痛症；亦會於下丘腦調節體溫的部位發揮作用，從而提供退熱效果；且不會刺激胃部。Phenylephrine Hydrochloride能迅速而長效舒緩感冒所引起的鼻塞。Caffeine是一種興奮劑，協助舒緩感冒所引起的疲勞，促進清醒，使您倍感舒適。維他命C能促進感冒迅速痊癒。Terpin Hydrate能增加支氣管分泌，是一種有效祛痰劑。Noscapine是一種中樞止咳藥物，能舒緩乾咳。

療效
舒緩傷風感冒的症狀：鼻塞、咽喉痛、發燒、咳嗽、頭痛、身體痛楚。

禁忌症
對此產品內任何成份敏感人士，切勿服用。
如正服用或在過去兩週期間曾服用單胺氧化酶抑制劑的人士，切勿服用此藥。
如你正在懷孕，切勿服用本藥。


注意事項
1. 在就診中倘有使用其他止痛、解熱或感冒藥劑者，避免使用「必理痛」傷風感冒特強。
2. 若病徵持續應立即請教醫生。
3. 請將此藥及其他藥品存放在兒童不易觸及的地方。
4. 請以下人士在服用本藥之前向醫生查詢：
a. 正服用Warfarin或抗凝血藥(如正服用抗凝血藥，並需要長期服用Acetaminophen，這可能會增加流血機會，但仍可以偶然地同時服用本產品)
b. 患有高血壓、心臟病、心血管病(如Raynaud's Phenomenon—當遇到寒冷或壓力的時候，手指或腳尖可能會感到疼痛)、糖尿病、過度活躍甲狀腺、青光眼、嗜鉻細胞瘤(腎上腺髓質腫瘤)、前列腺問題。
c. 正服用控制高血壓藥如beta-blocker。
d. 正服用增加或抑制胃口的藥物。
e. 正接受抗抑鬱的藥物治療如amitriptyline(其中一種三環抗抑鬱藥)。
f. 正服用心臟藥物如digoxin。
g. 腎、肝病患者。
h. 正在哺乳。請避免於哺乳期間服用此藥。

不良反應
服用此藥物可能引致的不良反應是罕有的，如頭痛、暈眩、焦慮、難以入睡、提高血壓、作嘔、嘔吐。
如服此藥後出現以下不良反應，請立即延醫診治：
a. 出現敏感徵狀，如皮膚出疹或瘙癢，間中還出現呼吸困難、嘴脣、舌頭、咽喉或面部腫脹。
b. 皮膚出疹、脫皮或口腔出現潰瘍。
c. 曾因服用阿司匹靈或其他非類固醇類抗炎藥出現呼吸困難的徵狀，及在服食本藥物期間出現相似徵狀。
d. 出現不能解釋的瘀傷或流血。
e. 失明(可能因眼內出現不正常高血壓所致)，這是十分罕見的(但青光眼患者可能較易有此反應)。
f. 不尋常的快速脈搏率又或者感到不尋常快或無規律的心跳。
g. 尿留症(排尿困難)，尤其是有肥大前列腺的男士。

Last Revised: Oct 2018
Hong Kong

CDP-904(HK)
910554

PHARMA CODE REF: IS-1358

Version: 5		Harmony AMS - Artwork Information Panel		
Manufacturing Site Number: 910554	Manufacturing Site(s): GSK_KUALA LUMPUR_MALAYSIA	Technical Drawing (Do NOT include version number): IMA_C80_INS_131X208_DRW		
Product Market Trade Name: Panadol Cold & Flu Extra	Approving Market(s): Hong Kong-HKG	Material Spec: (Do NOT include version number): MSP-1500032		
Print Process: Offset	Colour Standard Reference: For colour please refer to Industry Standard Pantone values	Material Type: Paper/Cardboard Leaflet paper	SUBSTRATE	
Total Colours & Varnishes: 1		Body Text Size: 5.0pt	Smallest Text Size: 5.0pt	
BLACK		Leading: 6.5pt		
Total Special Finishes: 0		Horizontal Scale: 100%		
		Microtext: N		
Additional Info (1): N/A	Additional Info (2): N/A	Additional Info (3): N/A		
Artwork copyright is the property of the GSK Group of Companies. All suppliers providing a service to GSK for printed components of any description must ensure that they have a licence for all fonts / software used in conjunction with GSK artwork. The distribution and use of fonts / software without a licence constitutes an intellectual property infringement. GSK will not accept any liability for the breach of third party intellectual property rights by printed component suppliers. The GSK certification / audit process requires suppliers to declare that they do not use unlicensed fonts / software and may require the supplier to produce evidence of such licence to GSK.				

200 mm Measuring Bar

If an e-banner DOES NOT appear on the top of this document, THEN this document has NOT been printed from the Harmony system.

IMPORTANT

GSK LOC is responsible to approve the change documentation, artwork brief and final artwork, ensuring that it is accurate, consistent and complete.

GSK SDC is responsible for site technical requirements and pre-press suitability.

GSK Market is responsible to advise SDC when changes required impact the following:

**Formulation
Tablet embossing
Storage conditions
Shelf Life**

NOTE TO MARKET

Local approvers must ensure that trade mark and copyright statements included in the brief comply with guidance provided by Legal: Global Trade Marks.