


 <p>S. González, s.l.</p> <p>Mejora de la salud y el bienestar 28970 Humanes (Madrid) T: +34 91 697 65 02 - F: 91 697 18 03 www.sgonzalez.com</p>	GLAXOSMITHKLINE Code: 11000000008345	Not approved <input type="checkbox"/> Approved <input type="checkbox"/>
	APPROVED By Leonard DiChiara at 11:31 pm, Oct 06, 2011	Description: PROSP PANADOL BLACK HONG KONG 1105 Size: 10x50 mm Folded: Not Folded No. Colours: 2/2 Colour: Black P/2593
Version No: 1 Date: 28.9.2011	Fonts [IMAGE] Design received in Illustrator Fonts outlined Diseño realizado a escala: 100%	







必理痛

TM

傷風+感冒

傷風感冒熱飲

效用：含有真正黑加侖子味及豐富維他命C，迅速舒緩傷風感冒引起的症狀：發燒、頭痛、四肢酸疼、鼻塞、咽喉痛、寒顫及促進感冒痊癒。

每包含有
Acetaminophen 600mg, Ascorbic Acid (Vitamin C) 40mg, Phenylephrine Hydrochloride 10mg。

服法
只供成人及十二歲或以上之兒童服用。
將一包「必理痛」傷風感冒熱飲放入杯內，注入半杯熱水，攪勻並加入適量開水，如有需要亦可加糖。每隔四小時服用一包，二十四小時內不可服用超過六包。不可服用多於指定份量。在沒有醫生指導下，請勿自行服用超過七日。

禁忌症
對此產品內任何成份敏感人士，切勿服用。如正服用或在過去兩週期間曾服用單胺氧化酶抑制劑的人士，切勿服用此藥。

注意事項

- 服用此藥時，請勿同時服用其他含Acetaminophen藥物、解充血藥及其他治療傷風感冒或咳嗽藥物，以免服用過量。
- 在懷孕或授乳期的婦女，服用此藥前需要徵詢醫生意見。
- 服用後如有暈眩感，請勿駕駛或操作機器。
- 服用後若症狀持續，請向醫生查詢。在沒有醫生指導下，請勿自行服用此藥超過七日。
- 如病人本身患有下列疾病，服用此藥前請先向醫生查詢：高血壓、心臟病、糖尿病、甲狀腺過度活躍症、青光眼、嗜鉻細胞瘤(腎上腺髓質腫瘤)、前列腺肥大、心血管病(如Raynaud's Phenomenon — 當遇到寒冷或壓力的時候，手指或腳尖可能會感到疼痛)、肝或腎病。
- 糖尿病患者在服用此藥時應注意，此藥每包(6克)含3.8克糖；另外，此藥含有phenylalanine，可能會對患有苯丙酮尿症的病人造成不良反應。
- 病人如正服用以下藥物，在服用此藥前請先向醫生查詢：
 - Warfarin或其他抗凝血藥(如正服用抗凝血藥，並需要長期服用Acetaminophen，這可能會增加流血機會，但仍可以偶然地同時服用本產品)。
 - 控制高血壓藥如beta-blocker。
 - 心臟病藥如digoxin。
 - 增加或抑壓胃口的藥物。


e. 抗抑鬱的藥物如Amiripityline(其中一種三環抗抑鬱藥)。

服藥過量
如根據服藥量指示使用，藥物中毒是極之罕有的。但若服用超過指定份量很多的Acetaminophen，可嚴重影響肝功能，其徵狀可能會延遲一星期後才出現。不過，在大部份情況，及早治理服用過量，是令肝功能回復正常。因此，若有服用過量的情況，即使無任何徵狀，亦應儘速治理。服用超過指定份量的Acetaminophen可以引起肝衰竭。

不良反應
服用此藥產生不良反應是十分罕見的。服用此藥您可能會出現以下情況：頭痛、暈眩、緊張、難以入睡、血壓增加、作嘔及嘔吐。
如服後出現以下徵狀，請停用本藥及立即延醫診治：
a. 出現敏感徵狀，如皮膚出疹或瘙癢，口中出現呼吸困難、喉嚨、舌頭、咽喉或面部腫脹。
b. 皮膚出疹、脫皮或口腔出現潰瘍。
c. 曾因服用阿司匹靈或其他非類固醇類消炎藥出現呼吸困難的徵狀，及在服食本藥物期間出現相似徵狀。
d. 出現不能解釋的瘀傷或流血。
e. 失明(可能因眼內出現不正常高血壓所致)，這是十分罕見的(但青光眼患者可能較易有此反應)。
f. 不尋常的快速脈搏率又或者感到不尋常快或無規律的心跳。
g. 尿原症(排尿困難)，尤其是有肥大前列腺的男士。

包裝
五包六克裝。

貯存方法
貯存於25°C以下及兒童不可觸及的地方。



11000000008345

S. González, s.l.
artes gráficas

Mercurio 13 - Pol. Ind. "Coherman"
28970 Humanes (Madrid)
T. 91 697 65 02 - F. 91 697 18 03
artigraficas@s-gonzalezsl.com

GLAXOSMITHKLINE	
Code: 11000000008345	
Description: PROSP PANADOL BLACK HONG KONG 1105	
Size: 100 x 150 mm	Folded: Not Folded
No. Colours: 2/2	Colour: Black P/2593
Pharmacode: 461	Paper: Offset white 50 g/m ²

Not approved	<input type="checkbox"/>
Approved	<input type="checkbox"/>
Date and Signed	

Version No: 1
Date: 28.9.2011

Fonts	[IMAGE] Design received in Illustrator Fonts outlined <i>Diseño realizado a escala: 100%</i>
--------------	--

Drawing
EMA/PROSP01 Ed.01

Reverso / Back

100 mm

Sense of Reading

30 mm 20 mm

COLD+FLU

COLD & FLU HOT REMEDY

FOR EFFECTIVE RELIEF OF COLD & FLU SYMPTOMS: FEVER, HEADACHE, ACHING LIMBS, BLOCKED NOSE, SORE THROAT, CHILLS AND HELP FOR FAST RECOVERY, WITH BLACKCURRANT TASTE AND VITAMIN C.

Each Sachet Contains
Acetaminophen 600mg, Ascorbic Acid (Vitamin C) 40mg, Phenylephrine Hydrochloride 10mg.

Dosage
Only for Adults and Children aged 12 years and over. Empty contents of sachet into a cup or mug. Half fill with hot water. Stir well. Add cold water as necessary and sugar if desired. One sachet can be taken every 4 hours, up to a maximum of 6 sachets in 24 hours. Do not exceed the stated dose. Please do not use more than 7 days without medical advice.

Contraindications
Panadol Cold & Flu Hot Remedy is contraindicated in patients with a previous history of hypersensitivity to any of the ingredients. For the patients who are taking, or have taken, within the last two weeks, monoamine oxidase inhibitors, please consult a physician before taking Panadol Cold & Flu Hot Remedy.

Precautions

- Please do not take this medicine with other Acetaminophen containing products, decongestant or cough & cold medicine to avoid overdose.
- Consult your physician before taking this medicine during pregnancy or breast feeding.
- Patients should be advised not to drive or operate machinery if affected by dizziness.
- If the symptoms persist, seek medical advice. Please do not use more than 7 days without medical advice.
- Medical advice should be sought before taking this medication in patients with these conditions: hypertension, cardiovascular disease, diabetes, hyperthyroidism; angle closure glaucoma, pheochromocytoma (a tumour near the kidney); enlargement of the prostate gland; occlusive vascular disease (e.g. Raynaud's Phenomenon, which may appear as pain in the fingers or toes in response to cold or stress), liver and kidney impairment.
- This product contains 3.8g sugar per dose. This should be taken into account in patients with diabetes. It also contains a source of phenylalanine. May be harmful for people with phenylketonuria.
- Please consult your doctor first if you are taking the following drugs:
 - Warfarin or similar medicines used to thin the blood (The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of Acetaminophen with increased risk of bleeding; occasional doses have no significant effect).
 - Medicines to control your blood pressure, such as beta-blockers.
 - Digoxin or similar medicines for heart disease.
 - Appetite suppressants or stimulants.
 - Medicines to treat depression such as tricyclic antidepressants e.g. amitriptyline.

Overdosage
Toxicity is extremely rare following administration of recommended dose. Nevertheless, massive overdose of Acetaminophen may adversely affect the liver of some people and signs or symptoms of overdose may be delayed for up to one week following ingestion. Nevertheless, liver damage is largely reversible if the patient is treated early, thus immediately medical management is required in the event of overdose, even if symptoms of overdose are not present, Acetaminophen overdose may cause liver failure.

Adverse Reactions
Undesirable effects are extremely rare. When using this product you may experience headache, dizziness, nervousness, difficulty sleeping, increased blood pressure, nausea and vomiting. Stop taking this medicine and consult your physician immediately if you have:

- Allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- A skin rash or peeling, or mouth ulcers.
- Previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction with this product.
- Unexplained bruising or bleeding.
- Loss of vision, which may be due to abnormally high blood pressure in the eye. This is very rare but is more likely to occur in those with glaucoma.
- An unusually fast pulse rate, or a sensation of an unusually fast or irregular heartbeat.
- Difficulty in passing water. This is more likely to occur in men with an enlarged prostate gland.

Pack Size
5 sachets of 6g.

Storage Condition
Store below 25°C and keep out of reach of children.

11000000008345

20 mm 30 mm

Sense of Reading



150 mm