

Panadol *Menstrual*

✓ For fast relief of period pain.

COMPACT 隨身裝

服法

成人：每次服一至二粒，需要時每隔四小時服用一次，二十四小時內不可服用超過八粒。兒童(十二歲以下)：請服用兒童「必理痛」退燒止痛藥丸或藥水。服用時避免飲用過量咖啡或茶。

Dosage

Adults: Take 1-2 caplets every 4 hours as required. Do not exceed 8 caplets in 24 hours. Children (under 12): Take Panadol for Children (in chewable tablets or suspension). Avoid drinking too much tea or coffee whilst taking these caplets.

注意：服食過量有危險。不可服用多於指定份量。請勿同時服用超過一種含撲熱息痛的藥物。不適當使用撲熱息痛可引致肝中毒及嚴重肝功能受損。請將此藥及其他藥品放在兒童不易觸及的地方。請貯存於25°C以下。

Caution. It is dangerous to exceed the stated dose. Do not exceed the stated dose. Do not take more than one Acetaminophen containing product.

Inappropriate use of Acetaminophen may potentially cause liver toxicity and severe liver damage. Keep this and all medications out of reach of children. Store below 25°C.

Each caplet contains Paracetamol (Acetaminophen) 500mg and Caffeine 65mg.

Manufactured by GlaxoSmithKline

(Dungarvan) Ltd., Knockbrack,
Dungarvan, Co. Waterford, Ireland.

Trade marks are owned by or licensed to the
GSK group of companies

© 2023 GSK group of companies or its licensor

Reg No.: HK-53827




(852) 3013 4435



hk.customer-relations@gsk.com

62000000203939



BN

EXP

必理痛

經痛配方

12 CAPLETS
粒



必理痛

經痛配方

主治：有效舒緩月經時
引起的各種痛症 ✓



12 CAPLETS
粒

gsk

主治：有效舒緩月經時引起的各種痛症。

禁忌症

對此產品的成份敏感人士，請勿服用。

注意事項

服用時請勿服用其他含Acetaminophen的藥物。請將此藥及其他藥品放置在兒童不易觸及的地方。若病徵持續，應請教醫生。腎、肝病患者，請於服用前向醫生查詢。不建議在授乳期或正在懷孕的婦女在沒有醫生指示下服用此藥。如正服用抗凝血藥（例如：Warfarin及其他Coumarins），並需要長期服用Acetaminophen，這可能會增加流血機會，但仍可偶然服用「必理痛」經痛配方。由於此藥物含有caffeine，應避免飲用過量含caffeine的飲品，如茶，咖啡，及其他含caffeine的飲料。吸收高份量的caffeine可以引致失眠，身體震動及心跳擺動所引起的不適。

不良反應

根據服量指示使用，「必理痛」經痛配方很少機會會引起嚴重的副作用及不會使腸胃不適；曾有報導罕有地敘述過皮膚過敏（如蕁麻疹）及支氣管痙攣惡化。如在服食此藥物期間出現以下敏感徵狀，請即停止服食本藥並應儘速延醫治理：曾因服用阿士匹靈或其他非類固醇類抗炎藥出現呼吸困難的徵狀，出現不能解釋的瘀傷或流血，皮膚出疹、脫皮或口腔出現潰瘍。

過量服藥

根據服量指示使用，藥物中毒是極之罕有的。但若服用超過指定份量之Acetaminophen，可嚴重影響肝功能，其徵狀可能會延遲一星期後才出現。不過，在大部份情況，及早治理服用過量，是可令肝功能回復正常的。因此，若有服用過量的情況，即使無任何徵狀，亦應儘速治理。服用超過指定份量的Acetaminophen可能會引起肝衰竭。

For fast effective relief of period pain.

Contraindications

Do not use if you are allergic to any of ingredients in this product.

Precautions

Do not use with other Acetaminophen-containing products. Keep this and all medications out of reach of children. If symptoms persist or you have been diagnosed with liver or kidney impairment, seek medical advice before taking this medication. Without medical advice, use during breast-feeding or pregnancy should be avoided. The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of Acetaminophen with increased risk of bleeding; occasional doses have no significant effect. This product contains caffeine. Avoid drinking too much tea or coffee whilst taking these caplets. High caffeine intake can result in difficulty sleeping and an uncomfortable feeling in the chest caused by fluttering heartbeat.

Adverse Reaction

When taken in recommended doses, Panadol Menstrual is unlikely to cause significant undesirable effects and will not upset the stomach. Skin reactions, such as urticaria, and aggravation of bronchospasm have been described rarely. Stop taking this medicine and seek medical advice immediately if you have an allergic reaction, have had shortness of breath in response to aspirin or NSAIDs, bruising, bleeding, skin rash/peeling or mouth ulcers. These reactions are rare.

Overdosage

Toxicity is extremely rare following administration of recommended dose. Nevertheless, massive overdose of Acetaminophen may adversely affect the liver of some people and signs or symptoms of overdose may be delayed for up to one week following ingestion. Nevertheless, liver damage is largely reversible if the patient is treated early, thus prompt medical attention is essential in all cases of suspected overdose even if there is no sign or symptom. Acetaminophen overdose may cause liver failure.

62000000203939