

195mm(H) X 140mm(W) X 20.5mm(D)
Panadol Shoulder pack

Panadol
PAIN RELIEF PATCH
(For Neck and Shoulder)



CONTAINS PAIN RELIEVING MEDICINAL INGREDIENTS
8 PATCHES



HALEON

必理痛



滲透止痛貼
肩頸剪裁，直擊疼痛


附送4片輔助貼，有需要時用
功效：肌肉及關節疼痛，如肩頸

8片

必理痛
滲透止痛貼 (肩頸適用)



含鎮痛藥用成分，8片



area for batch and
expiry date



滲透止痛貼 (肩頸適用)

必理痛



CONTAINS PAIN RELIEVING MEDICINAL INGREDIENTS
8 PATCHES



Panadol
PAIN RELIEF PATCH
(For Neck and Shoulder)

Panadol



紓緩不同部位
的肌肉疼痛：

肩、頸

腰背

手臂

肘、膝

Indications:

- Contains medicinal ingredients penetrating down into affected area and effectively soothe muscular and joint pain in various parts of body, e.g. shoulder, neck, waist, back, elbow and knee
- Water-base, low allergenic & gentle to skin
- Strong and immediate cooling effect and long lasting penetration to the affected area
- Highly stretchable, adhesive, soft, air permeable and easy to peel off

Packaging:

- 8 X 12cm per piece

Storage:

- Store below 30°C. Avoid direct sunlight
- Keep this and all medication out of reach of children

Precautions:

For external use only. This product should be applied only to intact, healthy skin and not to skin wounds or open injuries. This product should not come into contact with the eyes, nostrils, other mucous membranes and rash. In case of accidental contact, rinse well with water. Do not use with external heat, such as electric heating pad, as it may result in excessive skin irritations or skin burns. Some people may experience discomfort, particularly those with sensitive skin, if this product is used after a bath. If the condition does not improve or worsens within 7 days of starting treatment, patients should seek medical advice to exclude an alternative underlying cause of pain. Discontinue the treatment if excessive skin irritation occurs. Not for use in children under 12 years of age unless directed by a healthcare professional. Seizures may occur in children with a history of seizures. This product should not be applied on a skin surface near the nostrils of infants and children due to the presence of camphor. Caution should be used during concomitant administration with anticoagulants and antiplatelet medications as it may result in increased risk of bleeding. This product contains glycol salicylate and when applied to the skin, can be absorbed through the skin into the blood. Not recommended for use during pregnancy and breastfeeding. Consult your doctor immediately if you apply this product more than you should.

Side Effects:

- Application site hypersensitivity, application site reaction, skin burning sensation, rash and pruritus

Contraindications:

- Do not use if you are allergic to Acetylsalicylic acid or other non-steroidal anti-inflammatory drugs
- Contraindicated in patients with known hypersensitivity to glycol salicylate, menthol, camphor, tocopherol acetate or any other ingredient of the preparation

Dosage and Administration:

- Adults and children aged 12 and over: Apply no more than 3 times a day. Increase or decrease dosage according to the condition of the affected area. Use one patch at a time
- Open the pouch and take out a patch. Cut the patch to desired size. Peel off protective backing and apply sticky side to affected area
- Sweat and dirt may reduce the stickiness of the patch. Clean and dry the affected area to increase the stickiness. After application, wash your hand
- Use an adhesive plaster to fix the patch for easy peel off area such as joints

Active Ingredients (有效成分)

Glycol Salicylate	1.5%w/w
L-Menthol	1.0%w/w
DL-Camphor	0.5%w/w
DL- α -Tocopherol Acetate (Vitamin E)	0.3%w/w

Manufactured by Teh Seng Pharmaceutical Mfg. Co. Ltd. Secondary Factory, No.42-1, Huan Gong Road, Yong Kang District, Tainan City, Taiwan.
Trade marks are owned by or licensed to the Haleon group of companies.
©2025 Haleon group of companies or its licensor.
Reg. No. HK-56055

(852) 3013 4435
mystory.hk@haleon.com

component code: PR01176HKA



REF.: Pan Patch N&S 8s Apr25 w53M2
JOB: Pan Patch N&S 8s Rev
COL. USED: 4c
CLIENT: GSK
CONTACT: Wilson Wu
COUNTRY: H.K.
FOLLOW: klwai
1st CHECK: KL 14:10
2nd CHECK: KL 14:15
MODIFY DATE: 23-5-2025
STUDIO VERS.: 3
START DATE: 23-5-2023

WorkShop 5 Limited
design printing production
DISCLAIMER: Please check all artwork, text & color proof thoroughly. It is your responsibility as the client to check these points and accept full liability. All design & layout are the property of "Workshop 5 Ltd", it can not be used without their permission.
www.workshop5.com Tel: (852) 2544 5111 Fax: (852) 2545 0646

Die-cut CMYK

Panadol®


PAIN RELIEF PATCH

CONTAINS PAIN RELIEVING MEDICINAL INGREDIENTS
10 PATCHES



必理痛®

gsk



滲透止痛貼

即時清涼紓痛，持久滲透

附送輔助貼，易貼更穩固

功效：肌肉及關節疼痛，如腰、背、手肘和膝蓋

10片

必理痛®
滲透止痛貼

含鎮痛藥用成分，10片



area for batch and
expiry date

滲透止痛貼

必理痛®




CONTAINS PAIN RELIEVING MEDICINAL INGREDIENTS
10 PATCHES

Panadol®
PAIN RELIEF PATCH

Panadol®

紓緩不同部位的
肌肉疼痛：



功效：

- 含藥用成分，藥力深入患處，有效紓緩不同部位的肌肉及關節疼痛，如肩、頸、腰、背、手肘和膝蓋
- 水性基劑，不刺激皮膚，不容易過敏
- 即時清涼，持久滲透
- 彈性黏貼，柔軟透氣，易撕易貼，感覺舒適自然

包裝：

- 每片10 X 14cm

貯存：

- 請貯存於攝氏30度以下，避免陽光直射
- 請將此藥及其他藥品放置在兒童不易觸及的地方

注意事項：

- 只供外用
- 不適用於裂開或有開放性傷口的皮膚。避免與眼睛、黏膜或皮膚處接觸
- 請勿與熱敷墊同用
- 使用後若痛楚情況持續或惡化，請諮詢醫生意見
- 使用後若出現過度皮膚敏感，請停止使用
- 懷孕、授乳期間婦女及12歲以下兒童，使用前請諮詢醫生意見
- 如包裝損壞，請勿使用

禁忌症：

- 對 Acetylsalicylic acid 或其他非類固醇類抗炎藥有過敏反應者禁用本藥

用法：

- 成人及12歲或以上兒童：每天可換貼最多3次，依患部情況增加或減少劑量
- 打開鋁袋，取出止痛貼，把止痛貼切割成適當大小，剝去表面薄紙，貼於患部使用
- 患部有濕汗或污垢時會減少膏面之黏貼性，貼用前應先將患部清洗並抹乾以增強黏貼性
- 於關節等易脫落之部位，可使用輔助貼固定

另有必理絡特強滲透止痛貼及特強滲透止痛啫喱於有藥劑師的藥房發售。特強藥力，更迅速更有效。

Panaflex Max Pain Relief
Patch and Max Pain
Relief Gel are also
available in pharmacies.



Indications:

- Contains medicinal ingredients penetrating down into affected area and effectively soothe muscular and joint pain in various parts of body, e.g. shoulder, neck, waist, back, elbow and knee
- Water-base, low allergenic & gentle to skin
- Strong and immediate cooling effect and long lasting penetration to the affected area
- Highly stretchable, adhesive, soft, air permeable and easy to peel off

Packaging:

- 10 X 14cm per piece

Storage:

- Store below 30°C. Avoid direct sunlight
- Keep this and all medication out of reach of children

Precautions:

- For external use only
- Do not apply to torn skin or open wounds. Avoid contact with the eyes, mucous membranes or rashes
- Do not use with a heating pad
- Consult a doctor if symptoms persist or condition worsens
- Stop using if excessive skin irritation occurs
- Pregnant or nursing women, and children under 12 years of age: consult a doctor before use
- Do not use if the carton is broken

Contraindications:

- Do not use if you are allergic to Acetylsalicylic acid or other non-steroidal anti-inflammatory drugs

Dosage and Administration:

- Adults and children aged 12 and over: Apply no more than 3 times a day. Increase or decrease dosage according to the condition of the affected area
- Open the pouch and take out a patch. Cut the patch to desired size. Peel off protective backing and apply sticky side to affected area
- Sweat and dirt may reduce the stickiness of the patch. Clean and dry the affected area to increase the stickiness
- Use an adhesive plaster to fix the patch for easy peel off area such as joints

Active Ingredients (有效成分)

Glycol Salicylate	1.5%w/w
L-Menthol	1.0%w/w
DL-Camphor	0.5%w/w
DL- α -Tocopherol Acetate (Vitamin E)	0.3%w/w

Manufactured by Teh Seng Pharmaceutical Mfg. Co. Ltd.
Secondary Factory, No. 42-1, Hwan Gond Road, Yung
Kang City, Tainan Hsien, Taiwan, R.O.C.

必理痛，必理絡，Panadol and Panaflex
are trade marks owned by or licensed
to the GSK group of companies.
©2016 GSK or its licensor.
Reg. No: HK-56055

component code: PR-01168HKAB

152mm(H) X 213mm(W) X 29mm(D)

Die-cut CMYK



REF: Pan Patch GSL10s Apr16 S1M2
JOB: Pan Patch GSL10s
COL. USED:

CLIENT: GSK
CONTACT: gloria.h.tse@gsk.com

DATE: 20-4-2017
VERS.: 1
START: 21-4-2016

DISCLAIMER: Please check all artwork, text & color proof thoroughly. It is your responsibility as the client to check these points and accept full liability. All design & layout are the property of "WorkShop 5 Ltd." it can not be used without their permission.
www.workshop5.com Tel: (852) 2544 5111 Fax: (852) 2545 0646

Panadol[®]
PAIN RELIEF PATCH

CONTAINS PAIN RELIEVING MEDICINAL INGREDIENTS
18 PATCHES




必理痛[®]

gsk



area for batch and
expiry date



滲透止痛貼

即時清涼紓痛，持久滲透

附送輔助貼，易貼更穩固
功效：肌肉及關節疼痛，如腰、
背、手肘和膝蓋

18片

必理痛[®]
滲透止痛貼

含鎮痛藥用成分，18片




CONTAINS PAIN RELIEVING MEDICINAL INGREDIENTS
18 PATCHES

Panadol[®]
PAIN RELIEF PATCH

Panadol[®]

紓緩不同部位
的肌肉疼痛：



功效：

- 含藥用成分，藥力深入患處，有效紓緩不同部位的肌肉及關節疼痛，如肩、頸、腰、背、手肘和膝蓋
- 水性基劑，不刺激皮膚，不容易過敏
- 即時清涼，持久滲透
- 彈性黏貼，柔軟透氣，易撕易貼，感覺舒適自然

包裝：

- 每片10 X 14cm

貯存：

- 請貯存於攝氏30度以下，避免陽光直射
- 請將此藥及其他藥品放置在兒童不易觸及的地方

注意事項：

- 只供外用
- 不適用於裂開或有開放性傷口的皮膚。避免與眼睛、黏膜或皮膚處接觸
- 請勿與熱敷墊同用
- 使用後若痛楚情況持續或惡化，請諮詢醫生意見
- 使用後若出現過度皮膚敏感，請停止使用
- 懷孕、授乳期間婦女及12歲以下兒童，使用前請諮詢醫生意見
- 如包裝損壞，請勿使用

禁忌症：

- 對 Acetylsalicylic acid 或其他非類固醇類消炎藥有過敏反應者禁用本藥

用法：

- 成人及12歲或以上兒童：每天可換貼最多3次，依患部情況增加或減少劑量
- 打開鋁袋，取出止痛貼，把止痛貼切割成適當大小，剝去表面薄紙，貼於患部使用
- 患部有濕汗或污垢時會減少膏面之黏貼性，貼用前應先將患部清洗並抹乾以增強黏貼性
- 於關節等易脫落之部位，可使用輔助貼固定

另有必理痛特強滲透止痛貼及特強滲透止痛啫喱於有藥劑師的藥房發售。特強藥力，更迅速更有效。

Panaflex Max Pain Relief
Patch and Max Pain
Relief Gel are also
available in pharmacies.



Indications:

- Contains medicinal ingredients penetrating down into affected area and effectively soothe muscular and joint pain in various parts of body, e.g. shoulder, neck, waist, back, elbow and knee
- Water-base, low allergenic & gentle to skin
- Strong and immediate cooling effect and long lasting penetration to the affected area
- Highly stretchable, adhesive, soft, air permeable and easy to peel off

Packaging:

- 10 X 14cm per piece

Storage:

- Store below 30°C. Avoid direct sunlight
- Keep this and all medication out of reach of children

Precautions:

- For external use only
- Do not apply to torn skin or open wounds. Avoid contact with the eyes, mucous membranes or rashes
- Do not use with a heating pad
- Consult a doctor if symptoms persist or condition worsens
- Stop using if excessive skin irritation occurs
- Pregnant or nursing women, and children under 12 years of age: consult a doctor before use
- Do not use if the carton is broken

Contraindications:

- Do not use if you are allergic to Acetylsalicylic acid or other non-steroidal anti-inflammatory drugs

Dosage and Administration:

- Adults and children aged 12 and over: Apply no more than 3 times a day. Increase or decrease dosage according to the condition of the affected area
- Open the pouch and take out a patch. Cut the patch to desired size. Peel off protective backing and apply sticky side to affected area
- Sweat and dirt may reduce the stickiness of the patch. Clean and dry the affected area to increase the stickiness
- Use an adhesive plaster to fix the patch for easy peel off area such as joints

Active Ingredients (有效成分)

Glycol Salicylate	1.5%w/w
L-Menthol	1.0%w/w
DL-Camphor	0.5%w/w
DL- α -Tocopherol Acetate (Vitamin E)	0.3%w/w

Manufactured by Teh Seng Pharmaceutical Mfg. Co. Ltd.
Secondary Factory, No. 42-1, Hwan Gond Road, Yung
Kang City, Tainan Hsien, Taiwan, R.O.C.

必理痛，必理痛，Panadol and Panaflex
are trade marks owned by or licensed to
the GSK group of companies.
©2016 GSK or its licensor.
Reg. No: HK-56055

(852) 3013 4435
hk.customer-relations@gsk.com
www.gsk.com.hk

component code: PR-01177HKAB

152mm(H) X 213.5mm(W) X 41mm(D)

Die-cut CMYK



REF.: Pan Patch GSL18s Apr16 S1M2
JOB: Pan Patch GSL18s
COL. USED:
www.workshop5.com Tel.: (852) 2544 5111 Fax.: (852) 2545 0646

CLIENT: GSK
CONTACT: gloria.h.tse@gsk.com

DATE: 20-4-2017
VERS.: 1
START: 21-4-2016

DISCLAIMER: Please check all artwork, text & color proof thoroughly. It is your responsibility as the client to check these points and accept full liability. All design & layout are the property of "WorkShop 5 Ltd", it can not be used without their permission.

Panadol


PAIN RELIEF PATCH

CONTAINS PAIN RELIEVING MEDICINAL INGREDIENTS
36 PATCHES



HALEON

必理痛




滲透止痛貼

即時清涼紓痛，持久滲透

附送輔助貼，易貼更穩固
功效：肌肉及關節疼痛，如腰、背、手腳和膝蓋

36片

Pharmacode.29172



68mm

70mm

必理痛

滲透止痛貼

155mm(H) X 217mm(W) X 84mm(D)

Die-cut **CMYK**
280C

REF: Pan Patch GSL36s Apr25 w54M2 CLIENT: GSK FOLLOW: kiwai MODIFY DATE: 23-5-2025
JOB: Pan Patch GSL36s Rev CONTACT: Wilson Wu 1st CHECK: KL 12:10 STUDIO VERS.: 4
COL. USED: COUNTRY: H.K. 2nd CHECK: KL 12:15 START DATE: 1-6-2023

WorkShop 5 Limited
Creating Perfection Products

DISCLAIMER: Please check all artwork, text & color proof thoroughly. It is your responsibility as the client to check these points and accept full liability. All design & layout are the property of 'WorkShop 5 Ltd'; it can not be used without their permission.
www.workshops.com Tel: (852) 2544 5111 Fax: (852) 2545 0646

Panadol

舒緩不同部位
的肌肉疼痛：



功效：

- 含藥用成分，藥力深入患處，有效舒緩不同部位的肌肉及關節疼痛，如肩、頸、腰、背、手腳和膝蓋
- 水性基劑，不刺激皮膚，不容易過敏
- 即時清涼，持久滲透
- 彈性黏貼，柔軟透氣，易撕易貼，感覺舒適自然

包裝：

- 每片10 X 14cm

貯存：

- 請貯存於攝氏30度以下，避免陽光直射
- 請將此藥及其他藥品放置於兒童不易觸及的地方

注意事項：

只供外用。本產品只適用於完整，健康的皮膚。不適用於裂開或有開放性傷口的皮膚。避免與眼睛，鼻孔，其他黏膜和皮膚接觸。若有意外接觸，請用清水沖洗。請勿與外部熱能（例如電熱敷墊）同用，因有機會導致過度的皮膚刺激或灼傷。如在洗澡後使用，特別是有敏感皮膚的人士，可能會感到不適。若使用後的7日內，疼痛情況沒得到改善甚至惡化，病人應諮詢醫生意見，排除其他潛在疼痛原因。使用後若出現皮膚過度刺激，請停止使用。除非得到醫療專業人士建議，12歲或以下人士不適用。有腦癱發作紀錄的兒童有機會發作。因本產品含有樟腦，不應使用於有機會接觸嬰兒及兒童鼻孔附近的皮膚表面。注意與抗凝血和抗血小板藥物同用，因出血風險會增加。本產品含有 glycol salicylate，用於皮膚時能經皮膚被吸收到血液。不建議懷孕及授乳期間使用。如果使用此產品超過建議劑量，請立即諮詢醫生。

副作用：

- 用藥部位過敏，用藥部位反應，皮膚灼痛感，皮疹和瘙癢

禁忌症：

- 對 Acetylsalicylic acid 或其他非類固醇類抗炎藥有過敏反應者禁用本藥
- 對 glycol salicylate, menthol, camphor, tocopherol acetate 或本藥物其他成分有已知的過敏反應者禁用

用法：

- 成人及12歲或以上兒童：每天可換貼最多3次，依患部情況增加或減少劑量。每次使用一片止痛貼
- 打開鋁袋，取出止痛貼，把止痛貼切割成適當大小，剝去表面薄紙，貼於患部使用
- 患部有濕汗或污垢時會減少膏面之黏貼性，貼用前應先將患部清洗並抹乾以增強黏貼性。使用後，請洗手
- 於關節等易脫落之部位，可使用輔助貼固定

Indications:

- Contains medicinal ingredients penetrating down into affected area and effectively soothe muscular and joint pain in various parts of body, e.g. shoulder, neck, waist, back, elbow and knee
- Water-base, low allergenic & gentle to skin
- Strong and immediate cooling effect and long lasting penetration to the affected area
- Highly stretchable, adhesive, soft, air permeable and easy to peel off

Packaging:

- 10 X 14cm per piece

Storage:

- Store below 30 °C. Avoid direct sunlight
- Keep this and all medication out of reach of children

Precautions:

For external use only. This product should be applied only to intact, healthy skin and not to skin wounds or open injuries. This product should not come into contact with the eyes, nostrils, other mucous membranes and rash. In case of accidental contact, rinse well with water. Do not use with external heat, such as electric heating pad, as it may result in excessive skin irritations or skin burns. Some people may experience discomfort, particularly those with sensitive skin, if this product is used after a bath. If the condition does not improve or worsens within 7 days of starting treatment, patients should seek medical advice to exclude an alternative underlying cause of pain. Discontinue the treatment if excessive skin irritation occurs. Not for use in children under 12 years of age unless directed by a healthcare professional. Seizures may occur in children with a history of seizures. This product should not be applied on a skin surface near the nostrils of infants and children due to the presence of camphor. Caution should be used during concomitant administration with anticoagulants and antiplatelet medications as it may result in increased risk of bleeding. This product contains glycol salicylate and when applied to the skin, can be absorbed through the skin into the blood. Not recommended for use during pregnancy and breastfeeding. Consult your doctor immediately if you apply this product more than you should.

Side Effects:

- Application site hypersensitivity, application site reaction, skin burning sensation, rash and pruritus

Contraindications:

- Do not use if you are allergic to Acetylsalicylic acid or other non-steroidal anti-inflammatory drugs
- Contraindicated in patients with known hypersensitivity to glycol salicylate, menthol, camphor, tocopherol acetate or any other ingredient of the preparation

Dosage and Administration:

- Adults and children aged 12 and over: Apply no more than 3 times a day. Increase or decrease dosage according to the condition of the affected area. Use one patch at a time
- Open the pouch and take out a patch. Cut the patch to desired size. Peel off protective backing and apply sticky side to affected area
- Sweat and dirt may reduce the stickiness of the patch. Clean and dry the affected area to increase the stickiness. After application, wash your hand
- Use an adhesive plaster to fix the patch for easy peel off area such as joints

Active Ingredients (有效成分)

Glycol Salicylate	1.5%w/w
L-Menthol	1.0%w/w
DL-Camphor	0.5%w/w
DL-α-Tocopherol Acetate (Vitamin E)	0.3%w/w

Manufactured by Teh Seng Pharmaceutical Mfg. Co. Ltd.
Secondary Factory, No. 42-1, Huan Gong Road, Yong Kang District, Tainan City, Taiwan.

Trade marks are owned by or licensed to the Haleon group of companies.

©2025 Haleon group of companies or its licensor.

Reg. No. HK-56055

(852) 3013 4435
mystory.hk@haleon.com

component code: HK17C029-01



含鎮痛藥用成分，36片

必理痛

滲透止痛貼