


Miniature Pharma code Ref. No. 433



**Panadol Children's**  
*Used most by doctors*  
**FOR RELIEF OF FEVER AND PAINS, CONTAINS NO ASPIRIN**

**Description**  
For oral administration only.  
Each teaspoonful (5ml) of suspension contains Acetaminophen (Paracetamol) 120mg. Also contains Sucrose, Sorbitol Neosorb, Glycerin, Xanthan Gum, Strawberry Flavor, Methylparahydroxybenzoate, Carmoisine, Purified Water.

**Indications**  
Panadol for children reduces children's fevers quickly and effectively. Panadol for children also provides fast relief of mild-to-moderate pains associated with cold & flu, headaches, teething, immunizations.

**Contraindications**  
Panadol is contraindicated in patients with a previous history of hypersensitivity to any of the constituents.

**Precautions**  
This product contains paracetamol. Taking too much paracetamol can cause serious harm to your liver. Always read and follow the label. Keep this and all medicines out of reach of children. If symptoms persist or if you have been diagnosed with liver or kidney impairment, seek medical advice before taking this medication.  
**Please check with your doctor before use if you:**

- are underweight or malnourished
- regularly drink alcohol

You may need to avoid using this product altogether or limit the amount of paracetamol that you take.

- have a severe infection as this may increase the risk of metabolic acidosis.

Signs of metabolic acidosis include:

- deep, rapid, difficult breathing
- feeling sick (nausea), being sick (vomiting)
- loss of appetite

Contact a doctor immediately if you get a combination of these symptoms.

The anticoagulant effect of Warfarin and other Coumarins may be enhanced by prolonged regular daily use of Acetaminophen with increased risk of bleeding; occasional doses have no significant effect. Before taking this medicine, make sure you consult your doctor if you are taking warfarin or similar medicines used to thin the blood.

This medicine contains sorbitol liquid, patients with rare hereditary problems of fructose intolerance (deficiency in fructose metabolism) should not take this medicine.

Do not use this medicine if you are taking any other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu, or to aid sleep.

**Overdosage**  
Toxicity is extremely rare following administration of recommended dose. Nevertheless, massive overdose of Acetaminophen

may adversely affect the liver of some children and signs or symptoms of overdose may be delayed for up to one week following ingestion. Nevertheless, liver damage is largely reversible if the patient is treated early, thus prompt medical attention is essential in all cases of suspected overdose even if there is no sign or symptom. Administration of N-acetylcysteine or Methionine may be required. Acetaminophen overdose may cause liver failure.

**Adverse Reaction**  
Stop taking this medicine and seek medical advice immediately if you have:

- Allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- A skin rash or peeling, or mouth ulcers.
- Previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction with this product.
- Unexplained bruising or bleeding.

Undesirable effects are rare such as:

- Thrombocytopenia
- Anaphylaxis
- Cutaneous hypersensitivity reactions including skin rashes, angioedema, and Stevens Johnson Syndrome
- Bronchospasm in patients sensitive to aspirin and other NSAIDs
- Hepatic dysfunction


**Dosage and Administration**  
Suspension (Shake the bottle thoroughly before use).  
1 teaspoon is equivalent to 5ml. Each 5ml suspension (1 teaspoon) contains Acetaminophen 120mg.  
Panadol Suspension for Children is suitable for children aged 3 months and over. Not recommended in children under 3 months.  
To be taken at every 4 hours, or as prescribed by the physician.  
Dosage is calculated according to body weight of the patient:  
10-15mg/kg/dose (Acetaminophen/body weight/dose), i.e. dose of 10 to 15mg Acetaminophen for every kg of body weight.  
10mg/kg; weight of child (kg) x 10mg/kg = 24mg/ml.  
15mg/kg; weight of child (kg) x 15mg/kg = 24mg/ml.

Weight	Approximate dose (ml)
4 – 5.5kg (8.8 – 12.1 lbs)	2.5
5.5 – 7kg (12.1 – 15.4 lbs)	3
7 – 8.5kg (15.4 – 18.7 lbs)	4
8.5 – 11kg (18.7 – 24.2 lbs)	5
11 – 13.5kg (24.2 – 29.7 lbs)	6
13.5 – 16kg (29.7 – 35.2 lbs)	7
16 – 21kg (35.2 – 46.2 lbs)	9
21 – 27kg (46.2 – 59.4 lbs)	12
27 – 34kg (59.4 – 74.8 lbs)	15
34 – 42kg (74.8 – 92.4 lbs)	18

Do not take more frequently than every 4 hours.  
Do not exceed the stated dose. Do not take more than 4 doses in 24 hours.  
Maximum duration of continued use without medical advice: 3 days.  
Always use the lowest effective dose to relieve your symptoms.

**Storage conditions**  
Store below 25°C.  
Trade marks are owned by or licensed to the GSK group of companies.  
© 2017 GSK group of companies or its licensor

Miniature Pharma code Ref. No. 433



**必理痛 兒童**  
*最為醫學界採用*  
**主治：發燒、痛症。不含阿士匹靈。**

**成份**  
只供口服。  
每茶匙退燒止痛藥水 (5毫升) 含有Acetaminophen (Paracetamol) 120mg。亦含有Sucrose, Sorbitol Neosorb, Glycerin, Xanthan Gum, Strawberry Flavor, Methylparahydroxybenzoate, Carmoisine, Purified Water。

**療效**  
退燒、止痛。  
兒童必理痛能迅速有效地退燒，並能舒緩頭痛及因傷風感冒、出牙期間、疫苗注射所引起輕微至中等的痛症。

**禁忌症**  
對此產品內任何成分敏感人士，請勿服用。

**注意事項**  
本產品含撲熱息痛。若服用超過指定份量之撲熱息痛，可嚴重影響肝功能。請將此藥及其他藥品放置在兒童不易觸及的地方。請參閱及依照標籤指示使用本產品。服用後若病徵持續，應立即請教醫生。  
腎、肝病者，請於服用前向醫生查詢。如有以下狀況，在服用此藥前請先諮詢醫生：

- 體重過輕或營養不良
- 有飲酒習慣

避免與本產品一起使用或限制使用撲熱息痛的份量。

- 你有嚴重感染，嚴重感染會增加代謝性酸中毒的風險

代謝性酸中毒的徵狀包括：

- 呼吸深、急促及困難
- 感到不適 (噁心)；身體不適 (嘔吐)
- 食欲不振

如出現多於一個以上的徵狀，請立即求醫。

長期服用Acetaminophen，若同時服用抗凝血藥 (例如：Warfarin及其他Coumarins)，這可能會增加流血機會。但若偶然地同時服用兒童必理痛退燒止痛藥水及抗凝血藥，並無重要影響。如正服用Warfarin或抗凝血藥，請於服用此藥前向醫生查詢。

此藥水含有sorbitol，患有遺傳性果糖不耐症(果糖代謝缺陷)的人士，請勿服用此藥水。

如正服用處方或非處方撲熱息痛藥物以治療痛症、發燒、傷風感冒症狀、或用以幫助睡眠，請勿服用此藥。

**藥物過量**  
根據標籤指示服用，藥物中毒是極罕有的。但若服用超過指定份量很多的Acetaminophen，可嚴重影響肝功能，其徵狀可能會延遲一週後才出現。不過，在大部份情況，及早治療服用過量，是可令肝功能回復正常。因此，當有懷疑服用過量的情況，即使無任何徵狀，亦應儘速治療。可能需要以N-acetylcysteine或Methionine作治療。服用超過指定份量的Acetaminophen可能會引起肝衰竭。

**不良反應**  
如在服用此藥物期間出現以下徵狀，請即停止服用本藥並應儘速延醫治療：

- 出現敏感徵狀，如皮膚出疹或瘙癢，間中出現呼吸困難，喉嚨、舌頭、咽喉或面部腫脹
- 皮膚出疹、喉嚨或口腔出現瘙癢
- 曾因服用阿士匹靈或其他非類固醇類抗發炎藥出現呼吸困難的徵狀，及在服用本藥物期間出現相似徵狀
- 出現不能解釋的瘀傷或流血

不良反應是罕有的，如：

- 血小板減少症(Thrombocytopenia)
- 全身過敏性反應(Anaphylaxis)
- 皮膚敏感如出疹、血管性水腫(Angioedema)，由Stevens Johnson Syndrome引致的全身性多型性紅斑
- 對阿士匹靈及其他非類固醇類抗發炎藥敏感的病患者或曾出現支氣管痙攣(Bronchospasm)
- 肝機能障礙

**用法與劑量**  
藥水(服用前，請把退燒止痛藥水搖勻)。  
5毫升為1茶匙。每5毫升藥水(1茶匙)含有120毫克Acetaminophen。  
兒童必理痛退燒止痛藥水適合三個月或以上的兒童服用。不建議三個月以下嬰兒服用此藥。  
每四小時服用一次，或按醫生指示服用。  
服用劑量是按兒童本身的體重而計算的：  
每次10 – 15毫克/公斤(Acetaminophen/體重)，即每1公斤體重每次服用10至15毫克Acetaminophen。  
10毫克/公斤：兒童體重(公斤) x 10毫克/公斤 = 24毫克/毫升。  
15毫克/公斤：兒童體重(公斤) x 15毫克/公斤 = 24毫克/毫升。

體重	大約份量 (毫升)
4 – 5.5公斤 (8.8 – 12.1磅)	2.5
5.5 – 7公斤 (12.1 – 15.4磅)	3
7 – 8.5公斤 (15.4 – 18.7磅)	4
8.5 – 11公斤 (18.7 – 24.2磅)	5
11 – 13.5公斤 (24.2 – 29.7磅)	6
13.5 – 16公斤 (29.7 – 35.2磅)	7
16 – 21公斤 (35.2 – 46.2磅)	9
21 – 27公斤 (46.2 – 59.4磅)	12
27 – 34公斤 (59.4 – 74.8磅)	15
34 – 42公斤 (74.8 – 92.4磅)	18

請勿相離少於四小時服用此藥物。不可服用多於指定劑量。24小時內不可服用此藥多過4次。  
在沒有醫生指導下，不可持續使用超過三天。請使用最低有效劑量，以減輕症狀。

**貯存方法**  
貯存於25°C以下。  
將藥水儲藏於避光處或經授權使用。  
© 2017 葛蘭素史克集團或其授權人。

TEXT SIZE CONTAINED IN THIS ARTWORK		
Body text size: 6.0pt	Leading: 6.0pt	Horizontal Scale: 100%
Smallest text size: 6.0pt		Microtext: N

**FONT REPLACED**

Market to check  
all text 100%

<b>SGS Number:</b> 5008683		<b>Item Number:</b> GPI-PSSHK-2		<b>Manufacturing Site:</b> GSK-PHL-Cainta-PHCAI	
<b>GlaxoSmithKline Artwork Information Panel</b>		<b>Market Trade Name:</b> PANADOL Children's Suspension		<b>Colour Standard Reference Number:</b> N/A	
<b>Market or Pack Owner:</b> CH-Hong Kong- HKG		<b>Printing Process:</b> Offset		<b>Substrate:</b> White paper	
<b>Technical Reference No(s):</b> CAILFT0005_D_001 <small>(do NOT include the technical reference doc[s] version no[s].)</small>					
<b>Colours</b>	<b>K</b>				
<b>Varnishes</b>					
<b>Special Finishes</b>					
<b>Total: 1</b>		<b>Total: N/A</b>		<b>Total: N/A</b>	
				<b>RSC A/W Version: 5</b>	

Artwork copyright is the property of the GlaxoSmithKline Group of Companies  
All suppliers providing a service to GSK for printed components of any description must ensure that they have a licence for all fonts / software used in conjunction with GSK artwork. The distribution and use of fonts / software without a licence constitutes an intellectual property infringement. GSK will not accept any liability for the breach of third party intellectual property rights by printed component suppliers. The GSK certification / audit process requires suppliers to declare that they do not use unlicensed fonts / software and may require the supplier to produce evidence of such licence to GSK.

ATTENTION	ATTENTION	ATTENTION	ATTENTION	ATTENTION	ATTENTION	ATTENTION	ATTENTION
<b>To Ensure Accurate PDF Viewing and Printing:</b> FOR SCREEN VIEWING: Use Adobe Acrobat 7 Professional or Adobe Acrobat Reader, Standard or Professional (higher than 7). Overprint Preview must be activated for accurate on screen viewing. FOR PRINTING: Use only Acrobat Professional version 7 or higher. "Apply Overprint Preview" or "Simulate Overprinting" must be activated in the print settings for printing accurate hard copies.							
180 mm Measuring Bar							
If a status identification banner DOES NOT appear on this document, THEN this document has NOT been printed from the Global Pack Management system.							

**IMPORTANT**

GSK Market is responsible for this product, its design and content.  
Ensure the artwork is thoroughly checked, all the text proof-read and approved.  
RSC GSK is responsible for site technical requirements and pre-press suitability.  
GSK Market is responsible to advise RSC in case changes required impact the followings:  
**Formulation  
Tablet embossing  
Storage conditions  
Shelf Life**